

IMAGES IN CLINICAL RADIOLOGY

Laryngeal Trauma Complicating a Suicide Attempt by Hanging

Pierre Tritschler*, Gilles Delahaut* and Xavier Pavard*

A 40-year-old man was admitted to the emergency room after a suicide attempt. He was found hanged in his garden shelter by a neighbour. The patient complained only of a slight dysphagia and anterior cervical pain. The clinical examination revealed no cutaneous lesion or neurological deficit. Head and neck CT scan with and without injection of contrast agent was performed. CT has shown displaced fractures of the two horns of the thyroid cartilage (Figs. A, B; white arrow). Fractures were associated with a small tumefaction of the left hypopharyngeal soft tissue (Fig. C; white star) and left side of the larynx. The airways were patent. The ENT specialist achieved a laryngoscopy and confirmed the small bulging of the larynx, with



Figure A.



Figure B.

no significant effect on the airways. Larynx mobility was conserved. A control fibroscopy was done 12 hours later with no modification of the lesions.

Vascular structures were patent, without thrombosis or dissection. No bone lesion of cervical spine was found. Management consisted of conservative treatment and psychiatric support.

Comment

Fractures of the larynx are uncommon trauma that are associated with other life threatening injuries in about 45% of cases. The thyroid cartilage is the most often affected structure and horizontal tears are classically detected in strangulation cases. They are classically bilateral in the thyroid cartilage and affect the superior edge of the lamina and the superior horns [1]. These kind of

*Cliniques Universitaires Saint Luc Bruxelles, BE
 pierre.tritschler@hotmail.com, delahautgilles@hotmail.com,
 xavpavard@hotmail.com




Figure C.

fractures are often combined with hyoid bone tears and hypopharyngeal hematoma. 3D volume rendering is a good tool in association with axial CT images for the detection of these fractures. The diagnosis of fracture can be difficult because of the progressive ossification of the thyroid cartilage and differs from one person to another. The diagnosis is probably undervalued which can cause complications. Patient with nondisplaced laryngeal fractures can have long-term voice change if they are not treated. Butler et al reported that the voice results of a patient who were treated early after injury had a recovery rate that was more than two times than patients who have delayed treatment. Displaced thyroid and cricoid cartilage tears should be surgically reduced et fixed. To conclude, MDCT (2D and 3D VR) is crucial tool to detect quickly laryngeal injuries and other associated injuries.

Competing Interests

The authors declare that they have no competing interests.

Reference

1. **Becker, M, Leuchter, I, Platon, A, Becker, CD, Dulguerov, P and Varoquaux, A.** Imaging of laryngeal trauma. *Eur. J. Radiol.* Jan. 2014; 83(1): 142–154. DOI: <http://dx.doi.org/10.1016/j.ejrad.2013.10.021>. PMID: 24238937.

How to cite this article: Tritschler, P, Delahaut, G and Pavard, X 2016 Laryngeal Trauma Complicating a Suicide Attempt by Hanging. *Journal of the Belgian Society of Radiology*, 100(1): 2, pp. 1-2, DOI: <http://dx.doi.org/10.5334/jbr-btr.969>

Published: 28 January 2016

Copyright: © 2016 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.



Journal of the Belgian Society of Radiology is a peer-reviewed open access journal published by Ubiquity Press.

OPEN ACCESS 