

IMAGES IN CLINICAL RADIOLOGY

# Typical CT and MRI Features of Mucinous Rectal Adenocarcinoma


Jin-A Ryoo and Seung Soo Kim

**Main teaching point:** Mucinous rectal adenocarcinoma typically shows high signal intensity on T2-weighted images, weak enhancement, and internal calcification.


**Keywords:** Rectal neoplasms; Mucinous adenocarcinoma; Computed tomography; Magnetic resonance imaging

## Case History


A 69-year-old man was referred to our hospital for evaluation of a rectal mass that was incidentally detected at a local clinic. The patient was diagnosed with adenocarcinoma through colonoscopic biopsy, and computed tomography (CT) and magnetic resonance imaging (MRI) were performed for staging. Axial unenhanced and portal venous phase CT images (**Figure 1**) showed irregular and severe wall thickening (arrow and open arrow) in the rectum. The left side (open arrow) of the rectum contained calcification (arrowhead) and revealed poor enhancement relative to the right side (arrow). The majority of the wall thickening (open arrows) showed low and high signal intensity on T1- and T2-weighted imaging (**Figure 2A–C**), respectively. Gadolinium-enhanced MRI (**Figure 2D**) demonstrated greater enhancement of the right side rectal wall (arrow) compared with the left side wall (open arrow). On diffusion-weighted imaging ( $b = 800 \text{ s/mm}^2$ ) and apparent diffusion coefficient map (**Figure 3**), the well-enhanced right side rectal wall (arrow) revealed diffusion restriction, while there was no diffusion restriction in the left side wall (open arrow). The patient underwent Hartmann's operation after concurrent chemoradiotherapy, and the final diagnosis was mucinous rectal adenocarcinoma.



**Figure 1.**



**Figure 2.**



**Figure 3.**

## Comment

Mucinous rectal adenocarcinoma is a distinct pathologic subtype of rectal cancer. The World Health Organization defines mucinous adenocarcinoma as a neoplasm composed of greater than 50% extracellular mucin, which can be pathologically confirmed after surgical resection. Mucinous carcinoma has higher local recurrence, distant metastasis, lymph node metastasis, and venous invasion compared with nonmucinous carcinoma. Thus, a more aggressive surgical approach, including wide excision and extensive lymph node dissection, is needed in patients

with mucinous rectal adenocarcinoma than in patients with nonmucinous adenocarcinoma [1].

The abundant extracellular mucin of mucinous carcinoma results in typical CT and MRI features that are different from those observed in nonmucinous carcinoma. Mucinous carcinomas show less enhancement and more severe wall thickening in imaging compared with nonmucinous carcinomas. Moreover, calcium deposition within the wall thickening is the characteristic CT finding of mucinous adenocarcinoma. Extracellular mucin is hyperintense on T2-weighted images, and the presence of a T2-hyperintense signal greater than 50% of tumor volume suggests mucinous adenocarcinoma. Unlike nonmucinous components with diffusion restriction, the mucin components of mucinous adenocarcinoma do not exhibit diffusion restriction [1].

### Ethics and Consent

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amend-

ments or comparable ethical standards. This article does not contain any studies with animals performed by any of the authors.

Statement of informed consent was not applicable because the manuscript does not contain any patient data.

### Funding Information

This work was supported by the Soonchunhyang University Research Fund and the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (No. 2018R1C1B5085419).

### Competing Interests

The authors have no competing interests to declare.

### Reference

1. **Wnorowski, AM, Menias, CO, Pickhardt, PJ, Kim, DH, Hara, AK and Lubner, MG.** Mucin-containing rectal carcinomas: Overview of unique clinical and imaging features. *American Journal of Roentgenology*. 2019; 213(1): 26–34. DOI: <https://doi.org/10.2214/AJR.18.20864>

**How to cite this article:** Ryoo, J-A and Kim, SS. Typical CT and MRI Features of Mucinous Rectal Adenocarcinoma. *Journal of the Belgian Society of Radiology*. 2019; 103(1): 55, 1–2. DOI: <https://doi.org/10.5334/jbsr.1910>

**Submitted:** 19 August 2019

**Accepted:** 22 September 2019

**Published:** 04 October 2019

**Copyright:** © 2019 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.



*Journal of the Belgian Society of Radiology* is a peer-reviewed open access journal published by Ubiquity Press.

**OPEN ACCESS**